

Multiple & Mixed Quantifiers

Understanding Quantification

William Starr

11.01.11

Quantification

What We've Done

- ① So far, we've learned what \forall and \exists mean
 - Recall the **semantics** and **game rules**
 - Both based on **satisfaction**
- ② Use \forall and \exists for translation of quantifiers
 - Remember the **four Aristotelian Forms**
- ③ Two **logical concepts**
 - **FO Validity**
 - Logical truth restricted to $\forall, \exists, =, \neg, \wedge, \vee, \rightarrow, \leftrightarrow$
 - **FO Consequence**
 - Logical consequence restricted to $\forall, \exists, =, \neg, \wedge, \vee, \rightarrow, \leftrightarrow$
 - We test for these using the **replacement method**

Quantification

We are Just Getting Started

- This is a good start, but there is a lot more to understanding the logic of quantifiers
- Today we are going to think about what sentences containing **multiple quantifiers** mean
- As well as how to translate them into FOL
- We've only looked at sentences w/1 quantifier:
 - *All basketballs are orange*
 - *Some ninjas are not sociable*
- But what happens when there are 2, 3 or 4 quantifiers?

Quantification

Multiple Quantifiers

- Recall what old Abe said:

*You may fool **all** of the people **some** of the time; you can even fool **some** of the people **all** of the time; but you can't fool **all** of the people **all** of the time*
- Count the quantifiers: 6!
- The point:
 - We often communicate logically interesting things with several quantifiers
- So, as students of logic, we need learn how to mix multiple quantifiers

Multiple Existentials

A Simple Example

- We will begin by considering sentences with multiple occurrences of one quantifier

(1) *Some cube is left of some tetrahedron*

- How should we represent (1) in FOL?
- We have many options
- Let's consider and compare them

Multiple Existentials

Translating our Simple Example

(1) *Some cube is left of some tetrahedron*

- Two (of the many) correct translations:

(1a) $\exists x \exists y [\text{Cube}(x) \wedge \text{Tet}(y) \wedge \text{LeftOf}(x, y)]$

- There are objects x and y such that: x is a cube, y is a tetrahedron and x is left of y

(1b) $\exists x [\text{Cube}(x) \wedge \exists y (\text{Tet}(y) \wedge \text{LeftOf}(x, y))]$

- There is an object x such that x is a cube and there exists an object y such that y is a tetrahedron and x is left of y
- (1a) stacks all of the quantifiers at the beginning
 - This makes it easier to paraphrase
 - But less like the English (1)!

Multiple Existentials

Multiplicity of Translations

- In addition to:

(1a) $\exists x \exists y [\text{Cube}(x) \wedge \text{Tet}(y) \wedge \text{LeftOf}(x, y)]$

(1b) $\exists x [\text{Cube}(x) \wedge \exists y (\text{Tet}(y) \wedge \text{LeftOf}(x, y))]$

- We can put things in the reverse order:
 - (1c) $\exists y \exists x [\text{Cube}(x) \wedge \text{Tet}(y) \wedge \text{LeftOf}(x, y)]$
 - (1d) $\exists y [\text{Tet}(y) \wedge \exists x (\text{Cube}(x) \wedge \text{LeftOf}(x, y))]$
- Or put the predicates in a different order:
 - (1e) $\exists x \exists y [\text{Tet}(y) \wedge \text{Cube}(x) \wedge \text{LeftOf}(x, y)]$
 - (1f) $\exists x [\text{Cube}(x) \wedge \exists y (\text{LeftOf}(x, y) \wedge \text{Tet}(y))]$
- Let's look at these in Tarski's World to see that they are equivalent (`Equivalences.sen / .wld`)

Translation Convention

A Helpful Note

Translation Conventions (Stylistic Advice)

- All quantifiers are stacked up 'out in front'
- 1st quantifier in English sentence is written 1st and binds x , 2nd goes 2nd and binds y , etc.
- List predicates in order of quantifiers they restrict
 - Translate: *some cube is left of some tetrahedron*
 - (1a) $\exists x \exists y [\text{Cube}(x) \wedge \text{Tet}(y) \wedge \text{LeftOf}(x, y)]$
 - Rather than:
 - (1e) $\exists x \exists y [\text{Tet}(y) \wedge \text{Cube}(x) \wedge \text{LeftOf}(x, y)]$
 - $\text{Cube}(x)$ goes before $\text{Tet}(y)$ since $\exists x$ come before $\exists y$, $\text{Left}(x, y)$ goes last since it restricts neither $\exists x$ nor $\exists y$

Translation

Comments on Our Convention

- In general, there are very many different but equally correct ways of translating quantified sentences
 - Especially in sentences with multiple quantifiers
 - By equally correct we mean **FO Equivalent**
- Conventions on previous slide are **sylistic**
- Prenex Form**: all of a formula's quantifiers are stacked up at the front of the formula
 - Like: $\exists x \exists y (\text{Cube}(x) \wedge \text{Tet}(y))$
 - Not: $\exists y (\text{Cube}(x) \wedge \exists y \text{Tet}(y))$
- Everything we've said so far also holds for sentences containing multiple universal quantifiers

Multiple Universals

(2) *Every tetrahedron is larger than every cube*

- Given our conventions, the natural translation is:

(2a) $\forall x \forall y [(\text{Tet}(x) \wedge \text{Cube}(y)) \rightarrow \text{Larger}(x, y)]$

 - For every block x and every block y , if x is a tetrahedron and y is a cube then x is larger than y
- But this is equivalent to (among others):

(2b) $\forall x [\text{Tet}(x) \rightarrow \forall y (\text{Cube}(y) \rightarrow \text{Larger}(x, y))]$
- Let's look at Tarski's World
(`Equivalences.sen` / `.wld`)

Multiple Quantifiers

An Important Fact

Fact 1 (Multiplied Quantifiers)

When you have multiple occurrences of a single quantifier, **order does not matter**:

- $\exists x \exists y P(x, y) \Leftrightarrow \exists y \exists x P(x, y)$
- $\forall x \forall y P(x, y) \Leftrightarrow \forall y \forall x P(x, y)$

A Tricky Fact

Resisting the Temptation...

- It is tempting to paraphrase:

(3) $\forall x \forall y [(\text{Small}(x) \wedge \text{Cube}(y)) \rightarrow \text{RightOf}(x, y)]$

As:

(4) For every block x and every **other** block y , if x is small and y is a cube then x is right of y
- But **RESIST!**
 - (4) is **not** what (3) means
- (4) is really a paraphrase of:

(5) $\forall x \forall y [(x \neq y \wedge \text{Small}(x) \wedge \text{Cube}(y)) \rightarrow \text{RightOf}(x, y)]$
- (3) and (5) are **not equivalent**
- See this in TW (`Identity.sen`, `Identity.wld`)

The Tricky Fact

The Moral of the Story

The Tricky Fact

- 1 When evaluating sentences with multiple quantifiers, don't fall into the trap of thinking that distinct variables range over distinct objects
- 2 In fact, $\forall x \forall y P(x, y)$ logically entails $\forall x P(x, x)$, so the variables can't be assumed to range over distinct variables. (The same goes for \exists)

Mixing Quantifiers

Doing Things Differently



- In addition to repeating the same quantifier, you can mix quantifiers:
 - (6) *Everyone loves someone or other*
 - (7) *There is someone that everyone loves*
- Both (6) and (7) mix a universal and an existential
- But, they do it differently:
 - (6) is a *Universal Existential*
 - (7) is an *Existential Universal*
- Accordingly, we translate (6) and (7) differently:
 - (6') $\forall x \exists y (\text{Love}(x, y))$
 - (7') $\exists y \forall x (\text{Love}(x, y))$

Mixing Quantifiers

The Difference in Meaning is Big

- (6) *Everyone loves someone or other*
- (6') $\forall x \exists y (\text{Love}(x, y))$
- (7) *There is someone that everyone loves*
- (7') $\exists y \forall x (\text{Love}(x, y))$

- (6)/(6') and (7)/(7') describe **different** situations:



Mixing Quantifiers

Entailment Relations

- (6) *Everyone loves someone or other*
- (6') $\forall x \exists y (\text{Love}(x, y))$
- (7) *There is someone that everyone loves*
- (7') $\exists y \forall x (\text{Love}(x, y))$

Fact

(7) entails (6). By (7) there's some person, call him/her Pat, that everyone loves. It follows that everyone loves someone (or other), namely Pat!

Fact

(6) does **not** entail (7). Everyone could love a different person. Then (6) is true but (7) is not

Mixing Quantifiers

The Important Difference

- What examples (6) and (7) show is that when you mix quantifiers **order does matter!**
- This is very different from multiple occurrences of a single quantifier:
 - In that case, order does **not** matter
- To solidify the difference between *existential-universal* and *universal-existential* let's look at some examples in Tarski's World (MQ World.wld, MQ World 2.wld, MQ Sentences.sen)

Summary

Two Facts

Fact 1 (Multiplied Quantifiers)

When you have multiple occurrences of a single quantifier, **order does not matter:**

- 1 $\exists x \exists y P(x, y) \Leftrightarrow \exists y \exists x P(x, y)$
- 2 $\forall x \forall y P(x, y) \Leftrightarrow \forall y \forall x P(x, y)$

Fact 2 (Mixed Quantifiers)

When you have multiple occurrences of different quantifiers, **order does matter:**

- $\forall x \exists y P(x, y) \not\Leftrightarrow \exists y \forall x P(x, y)$

Exercise

Mixed Quantifiers in Tarski's World

- 11.11** (Building a world) Create a world in which all ten sentences in Arnault's Sentences are true.